



TITLE:

Some conjectures about the colored Jones polynomial (Intelligence of Low-dimensional Topology)

AUTHOR(S):

Tran, Anh T.

CITATION:

Tran, Anh T.. Some conjectures about the colored Jones polynomial (Intelligence of Low-dimensional Topology). 数理解析研究所講究録 2017, 2052: 74-81

ISSUE DATE:

2017-10

URL:

<http://hdl.handle.net/2433/237113>

RIGHT:

Some conjectures about the colored Jones polynomial

Anh T. Tran

Tohoku University/JSPS and University of Texas at Dallas

1 Introduction

The Jones polynomial was discovered by Jones in 1984 and has made a revolution in knot theory. For a knot K in S^3 , the colored Jones function $J_K(n)$ is a sequence

$$J_K : \mathbb{Z} \rightarrow \mathbb{C}[t^{\pm 1}]$$

of Laurent polynomials in t , which essentially measures the Jones polynomials of K and its parallels. Technically, $J_K(n)$ is the quantum invariant using the n -dimensional representation of sl_2 , normalized so that

$$J_{\text{unknot}}(n) = (t^{2n} - t^{-2n}) / (t^2 - t^{-2})$$

and extended to integers n by $J_K(n) = -J_K(-n)$.

In this note, we give a short survey about some old and new conjectures about the geometry and topology of the colored Jones polynomial. These includes:

- The volume conjecture of Kashaev [Ka] and Murakami–Murakami [MuMu] that relates the colored Jones polynomial of a knot in the 3-sphere and the hyperbolic volume of the knot complement.
- The AJ conjecture of Garoufalidis [Ga1] that relates the colored Jones polynomial and the A -polynomial of a knot. This conjecture is based on the theory of non-commutative A -polynomial of Frohman–Gelca–LoFaro [FGL] and the theory of q -holonomicity of quantum invariants of Garoufalidis–Le [GL].
- The slope conjecture of Garoufalidis [Ga3] and strong slope conjecture of Kalfagianni–Tran [KaTr] that relate the colored Jones polynomial of a knot and the boundary slopes and Euler characteristic of incompressible surfaces in the knot complement.

2 The volume conjecture

According to Thurston’s theory, by cutting the knot complement $S^3 \setminus K$ along appropriate disjoint tori one gets a collection of pieces, each is either Seifert fibered or hyperbolic; and $\text{Vol}(K)$ is defined as the sum of the hyperbolic volume of the hyperbolic pieces.

Then the volume conjecture of Kashaev [Ka] and Murakami–Murakami [MuMu] is stated as follows.

Volume conjecture. Suppose K is a knot in S^3 . Then

$$\lim_{n \rightarrow \infty} \frac{\log |J_K(n; t = e^{\pi i/2n})|}{n} = \frac{\text{Vol}(K)}{2\pi}.$$

The volume conjecture was confirmed for the following knots:

- 4_1 (by Ekholm),
- 5_2 (by Kashaev–Yokota [KY], Ohtsuki [Oh1]),
- torus knots (by Kashaev–Tirkkonen [KaTi]),
- Whitehead doubles of torus knots of type $(2, b)$ (by Zheng [Zh]),
- knots with up to 8 crossings (by Ohtsuki–Yokota [OY], Ohtsuki [Oh2], Takata [Ta]).

3 The AJ conjecture

Garoufalidis and Le [GL] proved that, for a fixed knot K , the colored Jones function $J_K(n)$ satisfies a non-trivial linear recurrence relation of the form

$$a_d(t^{2n}, t)J_K(n+d) + \cdots + a_0(t^{2n}, t)J_K(n) = 0$$

for all n , where $a_j(u, v) \in \mathbb{C}[u, v]$. For example, the colored Jones polynomial of the trefoil knot is

$$J_K(n) = t^{-6(n^2-1)} \sum_{j=-(n-1)/2}^{(n-1)/2} t^{24j^2+12j} \frac{t^{8j+2} - t^{-(8j+2)}}{t^2 - t^{-2}}.$$

It satisfies the following linear recurrence relation

$$(t^{8n+12} - 1)J_K(n+2) + (t^{-4n-6} - t^{-12n-10} - t^{8n+10} + t^{-2})J_K(n+1) - (t^{-4n+4} - t^{-12n-8})J_K(n) = 0.$$

Consider a discrete function $f : \mathbb{Z} \rightarrow \mathbb{C}[t^{\pm 1}]$, and define the linear operators M and L acting on such functions by:

$$(Mf)(n) = t^{2n}f(n), \quad (Lf)(n) = f(n+1).$$

Then $LM = t^2ML$. Moreover $M^{\pm 1}, L^{\pm 1}$ generate the *quantum torus* \mathcal{T} , a noncommutative ring with presentation

$$\mathcal{T} = \mathbb{C}[t^{\pm 1}] \langle M^{\pm 1}, L^{\pm 1} \rangle / (LM = t^2ML).$$

The *recurrence ideal* of the discrete function f is the left-ideal \mathcal{A} in \mathcal{T} annihilates f :

$$\mathcal{A} = \{P \in \mathcal{T} : Pf = 0\}.$$

Denote by \mathcal{A}_K the recurrence ideal of $J_K(n)$. Since $J_K(n)$ satisfies a non-trivial linear recurrence relation, $\mathcal{A}_K \neq 0$. The ring \mathcal{T} is not a principal left-ideal domain. However, by

adding the inverses of polynomials in t, M to \mathcal{T} , we get a principal left-ideal domain $\tilde{\mathcal{T}}$, and a generator α_K of the extension $\tilde{\mathcal{A}}_K = \tilde{\mathcal{T}}\mathcal{A}_K$. We call $\alpha_K \in \mathbb{C}[t, L, M]$ the *recurrence polynomial* of K . For example, the recurrence polynomial of the trefoil knot is

$$\begin{aligned} \alpha_K = & (t^{12}M^4 - t^{-4}M^{-2})L^2 \\ & + (t^{-6}M^{-2} - t^{-10}M^{-6}) - (t^{10}M^4 - t^{-2})L \\ & - t^4M^{-2} + t^{-8}M^{-6}. \end{aligned}$$

The AJ conjecture of Garoufalidis [Ga1] is stated as follows.

AJ conjecture. For any knot K in S^3 , $\alpha_K|_{t=-1}$ is equal to the A -polynomial of K , up to a factor depending on M only.

For example, for the trefoil knot

$$\alpha_K|_{t=-1} = (M^4 - 1)(L - 1)(LM^6 + 1)$$

is equal to the A -polynomial up to the polynomial $M^4 - 1$.

The A -polynomial of a knot in S^3 , defined by Cooper-Culler-Gillett-Long-Shalen [CCGLS], describes the $SL_2(\mathbb{C})$ -character variety of the knot complement as viewed from the boundary. It carries a lot of information about the knot. For example, the Newton polygon of the A -polynomial give rise to essential surfaces in the knot complement.

The AJ conjecture was confirmed for the following knots:

- $3_1, 4_1, 7_4$ (by Garoufalidis [Ga1], Garoufalidis–Koutschan [GK]),
- torus knots (by Hikami [Hi], Tran [Tr4]),
- some classes of two-bridge knots and pretzel knots (by Le [Le], Le–Tran [LT1], Le–Zhang [LZ]).

4 The slope conjecture

For a knot $K \subset S^3$, let X_K be its complement. Let $\langle \mu, \lambda \rangle$ be the canonical meridian–longitude basis of $H_1(\partial X_K)$. An element $p/q \in \mathbb{Q} \cup \{1/0\}$ is called a *boundary slope* of K if there is an essential surface $(S, \partial S) \subset (X_K, \partial X_K)$, such that ∂S represents $p\mu + q\lambda \in H_1(\partial X_K)$. We use bs_K to denote the set of boundary slopes of K . Hatcher [Ha] proved that bs_K is always a finite set.

Let $\hbar[J_K(n)]$ denote the highest degree of $J_K(n)$ in t . For every knot $K \subset S^3$, Garoufalidis [Ga3] proved that $\hbar[J_K(n)]$ is a *quasi-quadratic polynomial* in n . Namely, there exist periodic functions $a_K(n), b_K(n), c_K(n)$ such that

$$\hbar[J_K(n)] = a_K(n)n^2 + b_K(n)n + c_K(n).$$

Elements of $js_K := \{a(n)\}_n$ are called *Jones slopes* of K . Then the slope conjecture of Garoufalidis [Ga3] is stated as follows.

Slope Conjecture. For every knot $K \subset S^3$ we have

$$js_K \subset bs_K.$$

A similar conjecture can be stated for $\ell[J_K(n)]$, the lowest degree of $J_K(n)$ in t .

Example. For the $(-2, 3, 7)$ -pretzel knot we have

$$\begin{aligned}\hbar[J_K(n)] &= 37n^2/2 + 34n + e(n), \\ \ell[J_K(n)] &= 5n,\end{aligned}$$

where $e(n)$ is a periodic sequence of period 4. On the other hand, the $(-2, 3, 7)$ -pretzel knot is a Montesinos knot and its boundary slopes are given by $\{0, 16, 37/2, 20\}$. This confirms the slope conjecture for the $(-2, 3, 7)$ -pretzel knot.

The slope conjecture was confirmed for the following knots:

- alternating knots, knots with up to nine crossings, torus knots, $(-2, 3, 2n+1)$ -pretzel knots (by Garoufalidis [Ga3]),
- adequate knots (by Futer–Kalfagianni–Purcell [FKP]),
- 2-fusion knots (by Dunfield–Garoufalidis [DG], Garoufalidis–v.d.Veen [GV]),
- iterated cables of adequate knots (Kalfagianni–Tran [KaTr]),
- graph knots (Motegi–Takata [MT]),
- some classes of 3-stranded pretzel knots (Lee–v.d.Veen [LV]).

5 New conjectures about the colored Jones polynomial

Recall that for every knot $K \subset S^3$, there exist periodic functions $a_K(n), b_K(n), c_K(n)$ such that $\hbar[J_K(n)] = a_K(n)n^2 + b_K(n)n + c_K(n)$.

Then the following conjectures were proposed in [KaTr].

Conjecture 1 (Kalfagianni–Tran) For every non-trivial knot $K \subset S^3$, we have

$$b_K(n) \leq 0.$$

Note that $b_U(n) = 1/2$ for the trivial knot U .

Conjecture 2 [Strong slope conjecture] (Kalfagianni–Tran) Let K be a non-trivial knot and $r/s \in js_K$, with $s > 0$ and $\gcd(r, s) = 1$, a Jones slope of K . Then there is an essential surface $S \subset M_K$ with boundary slope r/s , and such that

$$\frac{\chi(S)}{|\partial S|s} \in \{b_K(n)\}_n.$$

Example. Consider the pretzel knot $K_p = (-2, 3, p)$, where $p \geq 5$ is an odd integer. Then, by [Ga3], we have

$$a_{K_p}(n) = \frac{p^2 - p - 5}{(p-3)/2} \quad \text{and} \quad b_{K_p}(n) = -\frac{(p-5)/2}{(p-3)/2}.$$

It is known [Ca] that K_p has an essential surface with boundary slope $\frac{p^2-p-5}{(p-3)/2}$, with 2 boundary components, and Euler characteristic

$$-(p-5) = 2\left(\frac{p-3}{2}\right)b_{K_p}(n).$$

This confirms the strong slope conjecture for the $(-2, 3, p)$ -pretzel knot.

Lee-v.d.Veen [LV] have recently verified the strong slope conjecture for some classes of 3-stranded pretzel knots.

6 Cable knots

In this last section, we discuss the above conjectures for cable knots. Suppose K is a knot, and p, q are coprime integers. The (p, q) -cable $K_{p,q}$ of K is the satellite of K with pattern (p, q) -torus knot. Then Morton [Mo] (see also [Ve]) proved that for $n > 0$ we have

$$J_{K_{p,q}}(n) = t^{-pq(n^2-1)/4} \sum_{k=-(n-1)/2}^{(n-1)/2} t^{4pk(qk+1)} J_K(2qk+1).$$

6.1 The volume conjecture for cable knots

Let E be the figure eight knot. By Habiro and Le, we have

$$J_E(n) = \frac{t^{2n} - t^{-2n}}{t^2 - t^{-2}} \sum_{k=0}^{n-1} \prod_{l=1}^k (t^{4n} + t^{-4n} - t^{4l} - t^{-4l}).$$

In this case we have the following result in [LT2].

Theorem. (Le–Tran) The volume conjecture holds true for the cable knot $E_{r,2}$ for every odd number r .

6.2 The AJ conjecture for cable knots

A formula for the A-polynomial of cable knots has recently been given by Ni–Zhang [NZ]. For an odd integer r we have

$$A_{K_{r,2}}(L, M) = \text{Res}_\lambda (A_K(\lambda, M^2), \lambda^2 - L) (M^{2r}L + 1).$$

On the other hand, from the cabling formula of Morton we have

$$J_{K_{r,2}}(n+1) = -t^{-2r(2n+1)} J_{K_{r,2}}(n) + t^{-2rn} J_K(2n+1).$$

Let $\mathbb{J}_K(n) := J_K(2n+1)$. Then, under certain conditions, we have

$$\alpha_{K_{r,2}} = \alpha_{\mathbb{J}_K} M^r (L + t^{-2r} M^{-2r}).$$

Using this, the AJ conjecture has been shown to hold true for

- most cable knots over torus knots (by Ruppe–Zhang [RZ]),
- most $(r, 2)$ -cable knots over the figure eight knot (by Ruppe [Ru], Tran [Tr2]), over some two-bridge knots (by Druivenga [Dr]).
- most $(r, 2)$ -cables of some classes of two-bridge knots and pretzel knots (by Tran [Tr3], [Tr1]).

6.3 The slope conjecture for cable knots

The following result relates the boundary slopes of a knot and its cable.

Theorem A (Motegi–Takata [MT]; Kalfagianni–Tran [KaTr]) For every knot $K \subset S^3$ and (p, q) coprime integers we have

$$(q^2 bs_K \cup \{pq\}) \subset bs_{K_{p,q}}.$$

By combining Theorem A and results about the degree of the colored Jones polynomial of adequate knots in [FKP], we have the following result in [KaTr].

Theorem B (Kalfagianni–Tran) Suppose K' is an iterated cable of an adequate knot K . Then K' satisfies the slope conjecture.

An iterated torus knot is an iterated cable of the trivial knot. Hence iterated torus knots satisfy the slope conjecture.

Motegi–Takata [MT] showed that the slope conjecture is preserved under connected sums. This, together with a modified version of Theorem B, implies that graph knots (knots whose hyperbolic volume is 0) satisfies the slope conjecture.

References

- [Ca] J. Callahan, *Denominators and differences of boundary slopes for $(1,1)$ -knots*, Bol. Soc. Mat. Mex. (3) **21** (2015), no. 2, 275–287.
- [CCGLS] D. Cooper, M. Culler, H. Gillet, D. Long and P. Shalen, *Plane curves associated to character varieties of 3-manifolds*, Invent. Math. **118** (1994), 47–84.
- [DG] Dunfield and S. Garoufalidis, *Incompressibility criteria for spun-normal surfaces*, Trans. Amer. Math. Soc. **364** (2012), no. 11, 6109–6137.
- [Dr] N. Druivenga, *The AJ-Conjecture for Cables of Two Bridge Knots*, preprint 2014, arXiv:1412.1053.
- [FGL] C. Frohman, R. Gelca, and W. Lofaro, *The A -polynomial from the noncommutative viewpoint*, Trans. Amer. Math. Soc. **354** (2002), 735–747.
- [FKP] D. Futer, E. Kalfagianni and J. Purcell, *Slopes and colored Jones polynomials of adequate knots*, Proc. Amer. Math. Soc. **139** (2011), 1889–1896.

- [Ga1] S. Garoufalidis, *On the characteristic and deformation varieties of a knot*, Proceedings of the Casson Fest, Geom. Topol. Monogr., vol. 7, Geom. Topol. Publ., Coventry, 2004, 291–309.
- [Ga2] S. Garoufalidis, *The degree of a q -holonomic sequence is a quadratic quasi-polynomial*, Electron. J. Combin. **18** (2011), no. 2, Paper 4, 23 pp.
- [Ga3] S. Garoufalidis, *The Jones slopes of a knot*, Quantum Topol. **2** (2011), no. 1, 43–69.
- [GK] C. Koutschan and S. Garoufalidis, *Irreducibility of q -difference operators and the knot 7_4* , Algebr. Geom. Topol. **13** (2013), no. 6, 3261–3286.
- [GL] S. Garoufalidis and T. Le, *The colored Jones function is q -holonomic*, Geom. Topol. **9** (2005), 1253–1293.
- [GV] S. Garoufalidis and R.v.d. Veen, *Quadratic integer programming and the slope conjecture*, New York J. Math. **22** (2016), 907–932.
- [Ha] A. Hatcher, *On the boundary curves of incompressible surfaces*, Pacific J. Math. **99** (1982), no. 2, 373–377.
- [Hi] K. Hikami, *Difference equation of the colored Jones polynomial for torus knot*, Internat. J. Math. **15** (2004), no. 9, 959–965.
- [Ka] R. Kashaev, *The hyperbolic volume of knots from the quantum dilogarithm*, Lett. Math. Phys. **39** (1997), no. 3, 269–275.
- [KaTr] E. Kalfagianni and A. Tran, *Knot cabling and the degree of the colored Jones polynomial*, New York J. Math. **21** (2015), 905–941.
- [KaTi] R. Kashaev and O. Tirkkonen, *A proof of the volume conjecture on torus knots*, J. Math. Sci. (N. Y.) **115** (2003), no. 1, 2033–2036.
- [KY] R. Kashaev and Y. Yokota, *On the volume conjecture for the knot 5_2* , preprint.
- [Le] T. Le, *The colored Jones polynomial and the A -polynomial of knots*, Adv. Math. **207** (2006), no. 2, 782–804.
- [LT1] T. Le and A. Tran, *On the AJ conjecture for knots*, Indiana Univ. Math. J. **64** (2015), no. 4, 1103–1151.
- [LT2] T. Le and A. Tran, *On the volume conjecture for cables of knots*, J. Knot Theory Ramifications **19** (2010), no. 12, 1673–1691.
- [LV] C. Lee and R.v.d. Veen, *Slopes for pretzel knots*, New York J. Math. **22** (2016), 1339–1364.
- [LZ] T. Le and X. Zhang, *Character varieties, A -polynomials, and the AJ conjecture*, Algebr. Geom. Topol. **17** (2017), no. 1, 157–188.
- [Mo] H. Morton, *The coloured Jones function and Alexander polynomial for torus knots*, Proc. Cambridge Philos. Soc. **117** (1995) 129–135.

- [MuMu] H. Murakami and J. Murakami, *The colored Jones polynomials and the simplicial volume of a knot*, Acta Math. **186** (2001), no. 1, 85–104.
- [MT] K. Motegi and T. Takata, *The slope conjecture for graph knots*, preprint 2015, arXiv:1501.01105.
- [NZ] Y. Ni and X. Zhang, *Detection of torus knots and a cabling formula for A -polynomials*, Algebr. Geom. Topol. **17** (2017), no. 1, 65–109.
- [Oh1] T. Ohtsuki, *On the asymptotic expansion of the Kashaev invariant of the 5_2 knot*, Quantum Topology **7** (2016), 669–735.
- [Oh2] T. Ohtsuki, *On the asymptotic expansion of the Kashaev invariant of the hyperbolic knots with 7 crossings*, preprint.
- [OY] T. Ohtsuki and Y. Yokota, *On the asymptotic expansions of the Kashaev invariant of the knots with 6 crossings*, preprint.
- [Ru] D. Ruppe, *The AJ conjecture and cabled knots over the figure eight knot*, Topology Appl. **188** (2015), 27–50.
- [RZ] D. Ruppe and X. Zhang, *The AJ conjecture and cabled knots over torus knots*, J. Knot Theory Ramifications **24** (2015), no. 9, 1550051, 24 pp.
- [Ta] T. Takata, *On the asymptotic expansions of the Kashaev invariant of some hyperbolic knots with 8 crossings*, preprint.
- [Tr1] A. Tran, *On the AJ conjecture for cable knots*, Fund. Math. **237** (2017), no. 3, 261–279.
- [Tr2] A. Tran, *On the AJ conjecture for cables of the figure eight knot*, New York J. Math. **20** (2014), 727–741.
- [Tr3] A. Tran, *On the AJ conjecture for cables of twist knots*, Fund. Math. **230** (2015), no. 3, 291–307.
- [Tr4] A. Tran, *Proof of a stronger version of the AJ conjecture for torus knots*, Algebr. Geom. Topol. **13** (2013), no. 1, 609–624.
- [Ve] R.v.d. Veen, *A cabling formula for the colored Jones polynomial*, preprint 2008, arXiv:0807.2679.
- [Zh] H. Zheng, *Proof of the volume conjecture for Whitehead doubles of a family of torus knots*, Chin. Ann. Math. Ser. B **28** (2007), no. 4, 375–388.

E-mail address: att140830@utdallas.edu